


BOLIVIA

Leishmaniasis cutaneous and mucosal 2015



Incidence
(100.000 pop.) **30.1**

Transmission
Leishmaniasis Composite Indicator (LCIc) **Moderate**

Occurrence

2231
NEW CASES

1995 cutaneous (89,4%)
236 mucosal (10,6%)

LCIc



AGE GROUP

5.1% < 5 years old
7.3% ≥ 5 < 10 years old
87.6% > 10 years old

12.4% of children
under 10 years old



SEX

69.3% Male
30.7% Female

44.2% of men
over 20 years old



CROSS-BORDER

Shares borders with
5 countries in **31**
provinces

43.7% of the cases
at the borders



SURVEILLANCE AND CONTROL

70.8% of the
population in
transmission areas

Individual Mandatory
Notification

No information on
vector chemical control



LEISHMANIA SPECIES

L. amazonensis
L. braziliensis
L. guyanensis
L. lainsoni



LUTZOMYIA VECTORS

L. flaviscutellata, *L.*
nuneztovari, *L. carrerai*
carrerai, *L. shawi*, *L.*
ayrozai, *L. yucumensis*, *L.*
llanosmartinsi, *L.*
nuneztovari anglesi



PATIENT ASSISTANCE

Free treatment at
healthcare facilities

Laboratory confirmation:
99.8%

Cure Rate: **99.7%**



AVAILABLE MEDICINES

No Information

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health -
National Leishmaniasis Programs of each country
Access on: December, 2016



Pan American
Health
Organization



World Health
Organization
REGIONAL OFFICE FOR THE
AMERICAS