

BOLIVIA

Cutaneous and mucosal Leishmaniasis 2017

Incidence
(100,000 pop.) **29.5**


Transmission
Triennium Leishmaniasis
Composite Indicator
(TLCIc 2015-2017) **Moderate**

Occurrence

2283
NEW CASES

2052 cutaneous (89.9%)

231 mucosal (10.1%)



**LCIc triennium
(2015-2017)**



AGE GROUP

6.6% < 5 Y/O
9.3% ≥ 5 < 10 Y/O
84.1% > 10 Y/O

15.9% children
under 10 years old



GENDER

68.6% Male
31.4% Female

60% men over 10
years old



CROSS-BORDER

Shares borders with
5 countries in **31**
provinces

25.3% of the cases
at the borders



SURVEILLANCE AND CONTROL

69.5% of the population
in transmission areas

Increase of the % of CL
in children under 10 Y/O
compared to the 2012-
2015 baseline (Targets of
the Plan of Action)



LEISHMANIA SPECIES

L. amazonensis
L. braziliensis
L. guyanensis
L. lainsoni



LUTZOMYIA VECTORS

L. flaviscutellata, *L.*
nuneztovari, *L. carrerai*
carrerai, *L. shawi*, *L.*
ayrozai, *L. yucumensis*,
L. llanosmartinsi, *L.*
nuneztovari anglesi



PATIENT ASSISTANCE

Laboratory confirmation:
100%

Cure rate: **58.6%**

% of cured patients among
the treated: **N/I**



AVAILABLE MEDICINE

Meglumine
Antimoniate

Liposomal
Amphotericin B

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health -
National Leishmaniasis Programs
Accessed on: 01 November 2018.

N/O: No information available; Y/O: years old