


PARAGUAY

Leishmaniasis cutaneous and mucosal 2015



LCIc



Incidence
(100.000 pop.) **7.93**

Transmission
Leishmaniasis
Composite Indicator
(LCIc) **Low**

Occurrence

126

NEW CASES

85 cutaneous (67.5%)

35 mucosal (27.8%)

06 not specified (4.7%)



AGE GROUP

2.4% < 5 years old
7.1% ≥ 5 < 10 years old
88.1% > 10 years old
2.4% not specified

9.5% of children under
10 years old



SEX

80% Male
20% Female

63.5% of men
over 20 years old



CROSS-BORDER

Shares borders with
3 countries in **23**
municipalities

23.8% of the cases
at the borders



SURVEILLANCE AND CONTROL

23,6% of the
population in
transmission areas

Individual Mandatory
Notification

Vector chemical control



LEISHMANIA SPECIES

L. braziliensis



LUTZOMYIA VECTORS

L. whitmani
L. neivai
L. migonei
L. intermedia



PATIENT ASSISTANCE

Free treatment at
healthcare facilities

Laboratory confirmation:
95.2%

Cure rate: **14.3%**



AVAILABLE MEDICINES

Meglumine antimoniate

Liposomal
Amphotericin B

Deoxycholate
Amphotericin B

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health -
National Leishmaniasis Programs of each country
Access on: December, 2016